

REC'D 1 2 JUL 2004

VIPO PCT

PA 1184198

# ARIO SER OUT REPORT OF THE CONTROL O

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

**United States Patent and Trademark Office** 

June 18, 2004

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.

APPLICATION NUMBER: 60/456,003

FILING DATE: March 19, 2003

PRIORITY DOCUMENT

COMPLIANCE WITH RULE 17.1(a) OP (b)

By Authority of the

COMMISSIONER OF PATENTS AND TRADEMARKS

P. SWAIN

**Certifying Officer** 

PTO/SB/16 (10-01)

Approved for use through10/31/2002. OMB 0651-0032

U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

# PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53(c). Express Mail Label No. EU639490479US

		IN	IVENTOR(	S)					
		Residence				7 60			
Given Name (first and middle [if	any])	Family Name of	or Sumame				ate or Fore	ign Country	) =
ROBERT VOIGT MULLEF				1	HEIDELBERG, GERMANY BIRKENAU, GERMANY				
CHRISTOPH CHRISTIAN		LUTZ		GIEBEN, GEI					
RUDI		SCHERHAG		SCHRIESHE					
Additional inventors are be	ing name	d on thi_1_sepai	rately numbe	ered sheets attac	hed he	reto		·	
	•	TITLE OF THE IN\	/ENTION (5	00 characters m	nax)				
CLOFARABINE DERIVATIVES				, •				•	• .,
				i i			1111111111111111		111)
Direct all correspondence to:		CORRESP	ONDENCE	ADDRESS					
K-71							ce Custon	e(VVnfger	( )
	Type Customer Number here Bar Code Lake Bar							FICE	
OR	Type Gus	tomer ivalliber nere							
Firm or Individual Name									
Address									
Address									
City			State			ZIP			
Country	1		Telephone			Fax			
Octavity	ENC	LOSED APPLICAT			apply		L		
Specification Number of		22		CD(s), Nu					
Drawing(s) Number of Si	hoete								
				Other (spe	ecify)	1			
Application Data Sheet. See 37 CFR 1.76									
METHOD OF PAYMENT OF FI	LING FEE	S FOR THIS PRO	VISIONAL A	PPLICATION FO	OR PA	TENT			
Applicant claims small e	entity statu	ıs. See 37 CFR 1,2	27					NG FEE	
A check or money orde	r is enclos	ed to cover the filir	ng fees				AML	OUNT (\$)	
The Commissioner is hereby authorized to charge filing 501786 \$80.00									
fees or credit any overp	-	•					9	30.00	
Payment by credit card.  The Invention was made by an				or under a cont	act with	h en ec	ency of the		
United States Government.	adenicy of	uia Offico States (		or unuor a contr	CIOL AALU	i aii ay	andy or all		
No.									
Yes, the name of the U.S. Gove	mment age	ency and the Governm	nent contract r	umber are:					
Respectfully submitted,		21 ml				1.11	03		
	S	/////////////	61-	Date	05	119/0	গ্ৰ		
SIGNATURE	12	The state of the s	1			RATION	NO.	44947	<u>'</u>
TYPED or PRINTED NAME STE	EPHEN J	MOLONEY /	/		approp	<i>riate)</i> umber:		1000.28.	000
	10 949 82			UC	NI IGNO	CITIDOI.	L	7000201	

#### TELEPHONE -USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT

This collection of information is required by 37 CFR 1.51. The information is used by the public to file (and by the PTO to process) a provisional application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 8 hours to complete, including gathering, preparing, and submitting the complete provisional application to the PTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, Washington, D.C. 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Box Provisional Application, Assistant Commissioner for Patents, Washington, D.C.

PTO/SB/17 (1-03)

Approved for use through 04/30/2003. OMB 0651-0032

Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

FEE IRANSMILIA	ᄔ			<u> </u>	T T T T T T T T T T T T T T T T T T T	ipiete ii Knowii			
for FY 2003			cation Nu	umber					
101 F 1 2003	Ļ	Filing Date			03/19/03	03/19/03			
Effective 01/01/2003. Patent fees are subject to annual revision	ON.	First Named Inventor N			VOIGT, ET AL.	VOIGT, ET AL.			
Applicant claims small entity status. See 37 CFR 1.2	27	Examiner Name							
		Group Art Unit							
TOTAL AMOUNT OF PAYMENT (\$) \$80	Attorney Docket No.			et No.	1000.28.000	1000.28.000			
METHOD OF DAVIDENT ALL STATES					CALCULATION /				
METHOD OF PAYMENT (check all that apply)	_				CALCULATION (co	ntinuea)			
Check Credit card Money Other None			NAL FEI Small En						
Deposit Account:	Fee	Fee	Fee Fe	8	Fee Description	on	Fee Paid		
Deposit Account 501786	Code 1051	(\$) 130	Code (\$ 2051		rcharge - late filing fee or	oath			
Number Deposit	1052	50	2052		rcharge - late provisional	filing fee or cover			
Account ILEX ONCOLOGY, INC.	1053	130	1053 1	she	eet n - English specification		===		
Name	ľ	2,520							
The Commissioner is authorized to: (check all that apply)	1804		1804 92		ling a request for ex parte reexamination esting publication of SIR prior to Examiner				
Charge (ee(s) Indicated below Credit any overpayments				acti	ion				
Charge any additional fee(s) during the pendency of this application	1805	51,840° 1805 1,840° Requesting publication of SIR after Examine action							
Charge fee(s) indicated below, except for the filing fee to the above-identified deposit account.	1251	110	2251	55 Ext	tension for reply within fir	st month			
FEE CALCULATION	1252	410	2252 2	205 Ext	tension for reply within se	cond month .			
1. BASIC FILING FEE	1253	930	2253 4	165 Ext	tension for reply within th	ird month			
Large Entity   Small Entity	1254	1,450	2254 7	725 Ext	tension for reply within fo	urth month			
Fee Fee Fee Fee Description Code (\$) Code (\$) Fee Paid	1255	1,970	2255 9	985 Ext	tension for reply within fif	th month			
1001 750 2001 375 Utility filing fee	1401	320	2401 1	160 No	tice of Appeal				
1002 330 2002 165 Design filing	1402	2 320	2402 1	160 Fili	ng a brief in support of a	n appeal			
1003 520 2003 260 Plant filing fee	1403	280	2403 1	140 Re	quest for oral hearing				
1004 750 2004 375 Reissue filing	1451	1,510	1451 1,5	510 Pet	tition to institute a public	use proceeding			
1005 160 2005 80 Provisional filing fee 80.00	1452	110	2452	55 Pet	tition to revive - unavoida	ible			
SUBTOTAL (1) (\$) \$80.00	1453	1,300	2453 €	550 Pei	tition to revive - unintenti	ona!			
2. EXTRA CLAIM FEES FOR UTILITY AND	1501	1,300	2501 6		ility issue fee (or reissue)	· · · · · · · · · · · · · · · · · · ·			
Fee from	1502				sign issue fee	· · · · · · · · · · · · · · · · · · ·			
Extra Claims below Fee Paid  Total Claims -20** = 0 X = 0.00	1503				ant issue fee				
Independent 3++= 0 X = 000	1460				titions to the Commission				
Ctains Muttiple Dependent	1807		1807	·	ocessing fee under 37 CF				
Large Entity   Small Entity	1806	180	1806 1		bmission of Information ( atement	Disclosure			
Fee Fee Fee Fee Description Code (\$) Code (\$)	8021	1 40	8021	40 Re	cording each patent assi				
1202 18 2202 9 Claims in excess of 20	1809	750	2809 3	•	nes number of properties Ing a submission after fin		<del></del>		
1201 84 2201 42 Independent claims in excess of 3				(37	7 ČFR § 1.129(a))				
1203 280 2203 140 Multiple dependent claim, if not paid	1810	750	2810 3	375 Foi (37	r each additional inventio 7 CFR § 1.129(b))	n io be examined			
1204 84 2204 42 ** Reissue independent claims	1801	750	2801 3		quest for Continued Example	mination (RCE)			
over original patent	1802	2 800	1802 - 9	900 Re	equest for expedited exam a design application	nination			
1205 18 2205 9 ** Reissue claims in excess of 20 and over original patent	C <sup>1</sup>	her lee (		Ol 1	a acsidii abbiicanoii				
SUBTOTAL (2) (\$) \$0.00		(					·		
SUBTOTAL (2) (\$) \$0.00	İ				CURTO	TAL (2)			
**or number previously paid, if greater, For Reissues, see above	'Re	educed b	y Basic Fil	ling Fee	Paid SUBTO	TAL (3) (\$)			
SUBMITTED BY					Complete	'if applicable)			
Name (Print/Type) STEPHEN J. MOLONE		Registration No. (Attorney/Agent) 44947				Telephone 210 949 8273			
~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	/_	(Allome)	yout			ļ			
Signature	W				Date	03/19/03			

WARNING: Information on this form may become public. Credit card information should not be included on this form. Provide credit card information and authorization on

This collection of information is required by 37 CFR 1.17 and 1.27. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 12 minutes to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the Individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Commissioner for Patents, Washington, DC 20231.

If you need assistance in completing the form, call 1-800-PTO-9199 (1-800-786-9199) and select option 2.

# PROVISIONAL APPLICATION COVER SHEET Additional Page

PTO/SB/16 (10-01)

Approved for use through 10/31/2002. OMB 0651-0032

U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a c ollection of information unless it displays a valid OMB control number.

INVENTOR(S)/APPLICANT(S)

Given Name (first and middle [if any])

DIETER

HERRMANN

HEIDELBERG, GERMANY

1000.28.000

Residence
(City, and either State or Foreign Country).

HEIDELBERG, GERMANY

Number 2 of 2

WARNING: Information on this form may become public. Credit card information should not be included on this form. Provide credit card information and authorization on PTO-2038.

# PROVISIONAL APPLICATION FOR UNITED STATES LETTERS PATENT

For

# **CLOFARABINE DERIVATIVES**

Ву

Robert Voigt, Christoph Müller, Christian Lutz, Rudi Scherhag, and Dieter Herrmann

## **BACKGROUND OF THE INVENTION**

The subject of the present invention are specific lipidesters of nucleotides of the general formula I,

#### wherein

- $R^1$  is a straight-chain or branched, saturated or unsaturated alkyl residue having 1-20 carbon atoms, optionally mono- or polysubstituted by halogen,  $C_1$ - $C_6$  alkoxy,  $C_1$ - $C_6$  alkylmercapto,  $C_1$ - $C_6$  alkoxycarbonyl,  $C_1$ - $C_6$  alkylsulfinyl or  $C_1$ - $C_6$  alkylsulfonyl groups,
- is hydrogen, a straight-chain or branched, saturated or unsaturated alkyl chain having 1-20 carbon atoms, optionally mono- or polysubstituted by halogen, C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>1</sub>-C<sub>6</sub> alkylmercapto, C<sub>1</sub>-C<sub>6</sub> alkoxycarbonyl or C<sub>1</sub>-C<sub>6</sub> alkylsulfonyl groups,
- R<sup>3</sup> represents hydrogen, hydroxy or halogen,
- R<sup>4</sup> represents hydroxy or hydrogen,
- R<sup>5</sup> represents hydrogen or hydroxy,
- R<sup>6</sup> represents a halogen,

with the proviso that at least one of the residues  $R^3$  or  $R^5$  is hydrogen, or  $R^3$  and  $R^4$  can signify a further bond between C-2' and C-3',

- X represents a valence bond, oxygen, sulfur, a sulfinyl or sulfonyl group,
- Y is a valence bond, an oxygen or sulfur atom, and
- n is an integer of 1, 2 or 3,

their tautomers and their physiologically acceptable salts of inorganic and organic acids and bases, as well as processes for their preparation and medicaments containing these compounds as active ingredients.

The amino group in the adenine residue of the general formula I can also be protected by well known amino protecting groups.

Since the compounds of the general formula I contain asymmetric carbon atoms, all optically-active forms and racemic mixtures of these compounds are also the subject of the present invention.

- J. Biol. Chem. 265, 6112 (1990) and EP-A-0,350,287 describe preparation and use of liponucleotides as anti-viral drugs. Therein, however, only dimyristoylphosphatidyl and dipalmitoylphosphatidyl residues coupled to well known nucleosides such as AZT and DDC are disclosed, including their fatty acid ester structure.
- J. Med. Chem. 33, 1380, (1990) describes nucleoside conjugates of thioether lipids with cytidine diphosphate, which have antitumor activity and might find use in oncology.

Chem. Pharm. Bull. 36, 209 (1988) describes 5'-(3-sn-phosphatidyl)nucleosides having antileukemic activity, as well as their enzymatic synthesis from the

corresponding nucleosides and phosphocholines in the presence of phospholipase D with transferase activity.

The patent application WO 92/03462 describes thioether lipid conjugates having antiviral activity, particularly for the treatment of HIV infections.

The synthesis of 2-Chloro-9-(2'-deoxy-2'-fluoro arabinofuranosyl)adenine (known as Clofarabine) is described in J. Org. Chem. 34, 2633 (1969) and in the patent application WO 01/60383.

The pharmacological activity of 2-Chloro-9-(2'-deoxy-2'-fluoro arabinofuranosyl)- adenine as inhibitor of DNA replication in comparison to other nucleosides is also described in Haematology 463 (1999).

Other halo arabinoadenosines with anticancer activity are mentioned in the patent applications US 5,384,310 and WO 92/20347.

The antiviral activity of such purine derivatives is shown in EP 0 314 011.

2-Chloro-9-(2'-deoxy-2'-fluoro arabinofuranosyl)adenine (Clofarabine) is a well known development product in clinical trials.

However, some toxic side effects of Clofarabine, mainly on the bone marrow, restrain the clinical use of this drug substance.

The compounds of the present invention in general formula I which incorporate the Clofarabine chemical structure, and homologues thereof, posses biological activity which distinguish them from the parent nucleosides. In particular, the compounds of the present invention show antitumoral activity and are useful in that at pharmacological relevant doses where one or more of the toxic side effects of the parent compound is/are ameliorated and/or the covalently

bound lipid part improves the bioavailability of the coupled drug substance and appears to contribute to enhanced selectivity and effectiveness of the compounds.

The compounds of the present invention have valuable pharmacological properties. In particular, they are suitable for therapy and prophylaxis of malignant tumors including, carcinomas, sarcomas, or leukemias.

Compared to the unconjugated nucleoside derivatives hitherto employed in treatment of malignant tumors, the compounds according to the invention have enhanced potency/efficacy or lower toxicity and thus, have a wider therapeutic window. In some embodiments of the present invention, the administration of pharmaceutical compositions comprising these compounds may be conducted continuously over a prolonged period of time. Incidences of withdrawal of the preparation or intermittent administration, which frequently are routine with chemotherapeutic agents due to their undesirable side-effects, may be reduced with the compounds according to this invention as compared to the parent compounds. Further, higher dose levels may be employed due to the amelioration of toxic side effects due to enhanced selectivity for tumor cytotoxicity.

The lecithin-like structure of the lipid moiety is desirable for the claimed improvements of the compounds of general formula I. The penetration through membranes and resorption barriers is facilitated and the conjugates according to formula I show a depository effect in different tissues.

The formation of lipid conjugates may also facilitate crossing the blood brain barrier due to better diffusion or active transport processes.

Similarly, the compounds of the present invention and their pharmaceutical formulations may be employed in free or fixed combination with other drugs for the treatment and prophylaxis of the diseases mentioned above.

Examples of these further drugs involve agents such as, e.g., mitosis inhibitors such as colchicines, vinblastine, alkylating cytostatic agents such as cyclophosphamide, melphalan, myleran or cis-platin, antimetabolites such as folic acid antagonists (methotrexate) and antagonists of purine and pyrimidine bases (mercaptopurine, 5-fluorouridine, cytarabine), cytostatically active antibiotics such as anthracyclines (e.g., doxorubicin, daunorubicin), hormones such as fosfestrol, taxanes, e.g. taxol, tamoxifen and other cytostatically/cytotoxically active chemotherapeutic and biologic agents.

Embodiments of the invention also encompass salts of the compounds of the general formula I, including alkali, alkaline earth and ammonium salts of the phosphate group. Examples of the alkali salts include lithium, sodium and potassium salts. Alkaline earth salts include magnesium and calcium and ammonium salts are understood to be those containing the ammonium ion, which may be substituted up to four times by alkyl residues having 1-4 carbon atoms, and/or aryl residues such as benzyl residues. In such cases, the substituents may be the same or different.

The compounds of general formula I may contain basic groups, particularly amino groups, which may be converted to acid addition salts by suitable inorganic or organic acids. To this end, possible as the acids are, in particular: hydrochloric acid, hydrobromic acid, sulphuric acid, phosphoric acid, fumaric acid, succinic acid, tartaric acid, citric acid, lactic acid, maleic acid or methanesulfonic acid.

In general formula I,  $R^1$  preferably represents a straight-chain  $C_8$ - $C_{16}$  alkyl residue which may be further substituted by a  $C_1$ - $C_6$  alkoxy or a  $C_1$ - $C_6$  alkylmercapto group.

More specifically, R<sup>1</sup> represents a nonyl, decyl, undecyl, dodecyl, tridecyl, tetradecyl or pentadecyl residue.

Preferably, methoxy, ethoxy, butoxy and hexyloxy groups are possible as substituents of  $R^1$  residue. In case  $R^1$  is substituted by a  $C_1$ - $C_6$  alkylmercapto residue, this is understood to be the methylmercapto, ethylmercapto, propylmercapto, butylmercapto and hexylmercapto residue, in particular.

Preferably,  $R^2$  represents a straight-chain  $C_8$ - $C_{15}$  alkyl group which may be further substituted by a  $C_1$ - $C_6$  alkoxy or a  $C_1$ - $C_6$  alkylmercapto group. More specifically,  $R^2$  represents an octyl, nonyl, decyl, undecyl, dodecyl, tridecyl or tetradecyl group. Preferably, methoxy, ethoxy, propoxy, butoxy and hexyloxy groups are preferable as the  $C_1$ - $C_6$  alkoxy substituents of  $R^2$ .

In case  $R^2$  is substituted by a  $C_1$ - $C_6$  alkylmercapto residue, this is understood to be the methylmercapto, ethylmercapto, propylmercapto, butylmercapto, pentylmercapto and hexylmercapto residue, in particular.

In some embodiments X is sulfur, sulfinyl or sulfonyl, and Y is oxygen.

Halogen is understood to be selected from the group of fluorine, chlorine, bromine, iodine, wherein fluorine and chlorine are preferred.

An example of a preferred lipid moiety is the group

#### wherein

 $R^1$  is  $C_{12}H_{25}$ 

 $R^2$  is  $C_{10}H_{21}$ 

X is S, SO or  $SO_2$ 

Y is 0.

Some embodiments include compounds wherein X and Y represent a valence bond,  $R^2$  is hydrogen, and  $R^1$  represents a  $C_1$ - $C_{20}$  alkyl chain optionally substituted by  $C_1$ - $C_6$  alkoxy or  $C_1$ - $C_6$  alkylmercapto.

In some embodiments, each  ${\bf R}^3$  and  ${\bf R}^4$  independently represent hydrogen or hydroxy.

In some embodiments, R<sup>5</sup> represents hydrogen.

In some embodiments, R<sup>6</sup> represents halogen, such as fluorine, chlorine or bromine, preferably fluorine.

In some preferred embodiments are compounds wherein  $R^3$  represents a hydroxy group,  $R^4$  and  $R^5$  are hydrogen and  $R^6$  is a halogen atom, such as fluorine.

In some embodiments n is 1.

Embodiments also include compounds of the general formula I were  $R^3$  and  $R^4$  together represent a further bond between C-2' and C-3'.

Preferred embodiments for the nucleoside residue in the general formula I include 2-Chloro-9-(2'-deoxy-2'-fluoro arabinofuranosyl)adenine and 2-Chloro-9-(arabinofuranosyl)adenine in their protected or unprotected form.

The compounds of the general formula I may be prepared by

1. reacting a compound of general formula II

R1-X-OP-OHOM (II)
$$0 - P - OHOM$$

wherein  $R^1$ ,  $R^2$ , n, X and Y have the meaning as indicated, with a compound of general formula III

wherein R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> and R<sup>6</sup> have the above-mentioned meaning, or represent a hydroxy group protected by an oxygen protecting group familiar to the artisan, in the presence of an activating acid chloride, such as 2,4,6-triisopropylbenzenesulfonic chloride, and a tertiary nitrogen base, e.g., pyridine or lutidine, in an inert solvent, such as toluene, or immediately in anhydrous pyridine, and optionally, subsequent to hydrolysis, removing the

oxygen protecting groups according to procedures conventional in nucleoside chemistry,

or

reacting a lipidalcohol (corresponding to formula II) with a nucleoside monophosphate (corresponding to formula III) in the same manner as mentioned above, or

2. reacting a compound of general formula IV,

wherein  $R^1, R^2$ , n, X and Y have the above-mentioned meaning, with a compound of general formula III,

wherein R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> and R<sup>6</sup> have the above-mentioned meaning, in the presence of phospholipase D from Streptomyces in an inert solvent, such as chloroform, in the presence of a suitable buffer, and optionally, subsequent to reaction, removing the oxygen protecting groups according to procedures conventional in nucleoside chemistry.

The preparation of the compounds of the general formula II and IV is performed in analogy to Lipids 22, 947 (1987) and J. Med. Chem. 34, 1377 (1991).

Compounds of formula III are prepared in analogy to J. Org. Chem. 34, 2633 (1969) and WO 01/60383.

Salts of compounds of general formula I are prepared by reacting the free acid with alkali or alkaline earth hydroxides, alcoholates or acetates.

The "enantiomers" in the lipid parts of the compounds of formula I may be prepared by separation via diastereomeric salts or by enantioselective synthesis of the lipid residues starting with optically active  $C_3$ -precursors.

The drugs containing compounds of formula I for the treatment of cancer may be administered in liquid or solid forms on the oral or parenteral route. Common application forms are possible, such as tablets, capsules, coated tablets, syrups, solutions, or suspensions.

Preferably, water is used as the injection medium, containing additives such as stabilizers, solubilizers and buffers as are common with injection solutions. Such additives are, e.g., tartrate and citrate buffers, ethanol, complexing agents such as ethylenediaminetetraacetic acid and its non-toxic salts, high-molecular polymers such as liquid polyethylene oxide for viscosity control. Liquid vehicles for injection solution need to be sterile and are filled in ampoules, preferably.

Solid carriers are, for example, starch, lactose, mannitol, methylcellulose, talc, highly dispersed silicic acids, higher-molecular fatty acids such as stearic acid, gelatine, agar-agar, calcium phosphate, magnesium stearate, animal and plant fats, solid high-molecular polymers such as polyethylene glycol, etc. If

desired, formulations suitable for oral application may include flavorings or sweeteners.

The dosage may depend on various factors such as mode of application, species, age, or individual condition.

The compounds according to the invention may suitably be administered orally or intravenously (i.v.) in amounts in the range of 0.1 - 100mg, preferably in the range of 0.2 - 80mg per kg of body weight and per day. In some dosage regimens, the daily dose is divided into 2-5 applications, with tablets having an active ingredient content in the range of 0.5 - 500mg being administered with each application.

Similarly, the tablets may have sustained release, reducing the number of applications, e.g., to 1-3 per day. The active ingredient content of sustained-release tablets may be in the range of 2-1000mg. The active ingredient may also be administered by i.v. bolus injection or continuous infusion, where amounts in the range of 5-1000mg per day are normally sufficient.

In addition to the compounds mentioned in the examples, the following compounds of formula I and their pharmacologically acceptable salts further exemplify compounds of the present invention:

- [2-Chloro-9-(2'-deoxy-2'-fluoro arabinofuranosyl)adenine]-5'phosphoric acid-(3-dodecylmercapto-2-decyloxy)propyl ester
- 2. [2-Chloro-9-(2'-deoxy-2'-fluoro arabinofuranosyl)adenine]-5'-phosphoric acid-(3-dodecylsulfinyl-2-decyloxy)propyl ester
- 3. [2-Chloro-9-(2'-deoxy-2'-fluoro arabinofuranosyl)adenine]-5'-phosphoric acid-(3-dodecylsulfonyl-2-decyloxy)propyl ester
- 4. [2-Chloro-9-(2'-deoxy-2'-fluoro arabinofuranosyl)adenine]-5'-phosphoric acid-(3-undecylmercapto-2-decyloxy)propyl ester

- 5. [2-Chloro-9-(2'-deoxy-2'-fluoro arabinofuranosyl)adenine]-5'-phosphoric acid-(3-undecylmercapto-2-undecyloxy)propyl ester
- 6. [2-Chloro-9-(2'-deoxy-2'-fluoro arabinofuranosyl)adenine]-5'-phosphoric acid-(3-decylmercapto-2-dodecyloxy)propyl ester
- 7. [2-Chloro-9-(2'-deoxy-2'-fluoro arabinofuranosyl)adenine]-5'-phosphoric acid-(3-dodecylmercapto-2-dodecyloxy)propyl ester
- 8. [2-Chloro-9-(2'-deoxy-2'-fluoro arabinofuranosyl)adenine]-5'-phosphoric acid-(3-decylmercapto-2-decyloxy)propyl ester
- [2-Chloro-9-(2'-deoxy-2'-fluoro arabinofuranosyl)adenine]-5'phosphoric acid-(3-undecylsulfinyl-2-decyloxy)propyl ester
- 10. [2-Chloro-9-(2'-deoxy-2'-fluoro arabinofuranosyl)adenine]-5'-phosphoric acid-(3-undecylsulfonyl-2-decyloxy)propyl ester
- 11. [2-Chloro-9-(2'-deoxy-2'-fluoro arabinofuranosyl)adenine]-5'-phosphoric acid-(3-undecylsulfinyl-2-undecyloxy)propyl ester
- 12. [2-Chloro-9-(2'-deoxy-2'-fluoro arabinofuranosyl)adenine]-5'-phosphoric acid-(3-undecylsulfonyl-2-undecyloxy)propyl ester
- 13. [2-Chloro-9-(2'-deoxy-2'-fluoro arabinofuranosyl)adenine]-5'-phosphoric acid-(3-tridecylmercapto-2-undecyloxy)propyl ester
- 14. [2-Chloro-9-(2'-deoxy-2'-fluoro arabinofuranosyl)adenine]-5'-phosphoric acid-(3-tridecylmercapto-2-decyloxy)propyl ester
- 15. [2-Chloro-9-(2'-deoxy-2'-fluoro arabinofuranosyl)adenine]-5'-phosphoric acid-(3-tridecylsulfinyl-2-decyloxy)propyl ester

## **EXAMPLE 1**

[2-Chloro-9-(2'-deoxy-2'-fluoro arabinofuranosyl)adenine]-5'-phosphoric acid-(3-dodecylmercapto-2-decyloxy)propyl ester

0.91 g of phosphoric acid-(3-dodecylmercapto-2-decyloxy)propyl ester are treated twice with 20 ml of anhydrous pyridine and concentrated by evaporation. The residue is dissolved in 20 ml of anhydrous pyridine at room

temperature, treated with 1.07 g of 2,4,6-triisopropylbenzenesulfonic chloride under nitrogen and stirred at 25°C for 0.5 hours. Then 0.5 g of 2-Chloro-9-(2'-deoxy-2'-fluoro arabinofuranosyl)adenine (Clofarabine) are added at once, and the charge is allowed to stand under nitrogen for 20 hours. Hydrolysis is performed by adding 5 ml of water, the mixture is stirred for another 0.5 hour at room temperature, freed from solvent under vacuum, and stripped twice using 50 ml of toluene. The residue is purified by column chromatography on Lichrospher 60 RPSelect B with methanol/aqueous 40mM sodium acetate 88:12 as the eluent. The product containing fractions are evaporated. The residue is distributed between 50 ml of tert.-butylmethylether and 10 ml of 2N hydrochloric acid. The organic layer is evaporated. The residue is dissolved in a mixture of 5 ml of toluene and of 5 ml of methanol. The pH is adjusted to pH 7 by addition of sodium methanolate. The solvent is stripped of and the residue is dried in vacuum.

The yield is 0.82 g (62%) white powder.

<sup>31</sup>P-NMR (121,5 MHz, D<sub>6</sub>-DMSO, 25°C): -0.4

Melting point: 176°C.

The phosphoric acid-(3-dodecylmercapto-2-decyloxy) propyl ester is prepared as described in WO 92/03462.

2-Chloro-9-(2'-deoxy-2'-fluoro arabinofuranosyl)adenine is prepared according to J. Org. Chem. 34, 2633 (1969) and WO 01/60383.

#### **CLAIMS**

## 1. A nucleotide derivative of formula l

$$R1 - X$$
 $R2 - Y$ 
 $O$ 
 $O$ 
 $R5$ 
 $R6$ 
 $O$ 
 $O$ 
 $R3$ 
 $R4$ 
 $R3$ 

#### wherein

 $R^1$  is selected from the group consisting of a straight-chain or branched, saturated or unsaturated alkyl chain having 1-20 carbon atoms, which is unsubstituted or substituted at least once by halogen,  $C_1$ - $C_6$  alkoxy,  $C_1$ - $C_6$  alkylmercapto,  $C_1$ - $C_6$  alkoxycarbonyl,  $C_1$ - $C_6$  alkylsulfinyl or  $C_1$ - $C_6$  alkylsulfonyl groups;

R  $^2$  is selected from the group consisting of hydrogen, a straight-chain or branched, saturated or unsaturated alkyl chain having 1-20 carbon atoms, which is unsubstituted or substituted at least once by halogen,  $C_1$ - $C_6$  alkylmercapto,  $C_1$ - $C_6$  alkoxycarbonyl or  $C_1$ - $C_6$  alkylsulfonyl groups;

R<sup>3</sup> is selected from the group consisting of hydrogen, hydroxy and halogen;

R<sup>4</sup> is selected from the group consisting of hydrogen and hydroxy;

R<sup>5</sup> is selected from the group consisting of hydrogen, hydroxy and halogen;

R<sup>6</sup> is halogen;

X is selected from the group consisting of a valence bond, an oxygen atom, a sulfur atom, a sulfinyl group and a sulfonyl group;

Y is selected from the group consisting of a valence bond, an oxygen atom and a sulfur atom;

n is an integer of 1, 2 or 3;

with the proviso that at least one of the residues R<sup>3</sup> or R<sup>5</sup> is hydrogen and that, furthermore, R<sup>3</sup> and R<sup>4</sup> can signify a further bond between C-2' and C-3', whereby the amino group of the nucleoside base may be unsubstituted or substituted by a known amino protecting group, their tautomers, their optically active forms and racemic mixtures, and their physiologically acceptable salts of inorganic and organic acids or bases.

- 2. The nucleotide derivative according to claim 1, wherein  $R^1$  is a straight-chain  $C_8$ - $C_{15}$  alkyl group, which is unsubstituted or substituted by a  $C_1$ - $C_6$  alkoxy or a  $C_1$ - $C_6$  alkylmercapto group.
- 3. The nucleotide derivative according to claim 1, wherein  $R^2$  represents a straight-chain  $C_8$ - $C_{15}$  alkyl group, which is unsubstituted or substituted by a  $C_1$ - $C_6$  alkoxy or a  $C_1$ - $C_6$  alkylmercapto group.
- 4. The nucleotide derivative according to claim 1, wherein X is sulfur, sulfinyl or sulfonyl, and Y is oxygen.
- 5. The nucleotide derivative according to claim 1, wherein X and Y are valence bonds,  $R^2$  is hydrogen, and  $R^1$  is a  $C_1$ - $C_{20}$  alkyl group, which is

unsubstituted or substituted by a  $C_1$ - $C_6$  alkoxy or a  $C_1$ - $C_6$  alkylmercapto group.

- 6. The nucleotide derivative according to claims 1 to 6, wherein n is 1.
- 7. The nucleotide derivative according to claims 1 to 6, wherein R<sup>5</sup> is selected from the group consisting of hydrogen, and halogen.
- 8. The nucleotide derivative according to claim 1, wherein R<sup>3</sup> and R<sup>4</sup> are individually selected from the group consisting of hydroxy or a hydrogen atom.
- 9. The nucleotide derivative according to claim 1, wherein R<sup>5</sup> is hydrogen.
- 10. The nucleotide derivative according to claim 1, wherein  ${\sf R}^6$  is fluoro.
- 11. The nucleotide derivative according to claim 1, wherein R<sup>3</sup> and R<sup>4</sup> signify a further bond between C-2' and C-3'.
- 12. The nucleotide derivative according to claim 1, wherein R<sup>4</sup> and R<sup>5</sup> are hydrogen.
- 13. The nucleotide derivative according to claim 12, wherein  ${\sf R}^{\sf 3}$  is hydroxy.
- 14. The nucleotide derivative according to claim 1, wherein

 $R^1$  is a straight-chain  $C_9$ - $C_{13}$  alkyl group which is unsubstituted or substituted by a methoxy, ethoxy, butoxy, hexyloxy, methylmercapto, ethylmercapto, propylmercapto, butylmercapto, or hexylmercapto residue;

 $R^2$  is a straight-chain  $C_8$ - $C_{12}$  alkyl group which is unsubstituted or substituted by a methoxy, ethoxy, butoxy, hexyloxy, methylmercapto, ethylmercapto, propylmercapto, butylmercapto, or hexylmercapto residue;

R3 is a hydroxy residue;

R4 is a hydrogen atom; and

R5 is a hydrogen atom.

- 15. The nucleotide derivative according to claim 1, wherein the nucleoside portion is 2-Chloro-9-(2'-deoxy-2'-fluoro arabinofuranosyl)adenine.
- 16. A composition comprising at least one compound according to claim 1 in combination with a pharmaceutically acceptable adjuvant or vehicle.
- 17. A method for treating malignant tumors comprising administering to a patient in need of such treatment an amount of the composition according to claim 16 effective to treat said tumors.
- 18. The method according to claim 17, wherein said tumor is selected from the group consisting of carcinomas, sarcomas or leukemias.
- 19. A method for treating malignant tumors comprising administering to a patient in need of such treatment an amount of the composition according to claim 16 effective to treat said tumors in fixed or free combination with other anticancer agents.

### 20. A nucleotide derivative of formula II

#### wherein

 $R^1$  is selected from the group consisting of a straight-chain or branched, saturated or unsaturated alkyl chain having 1-20 carbon atoms, which is unsubstituted or substituted at least once by halogen,  $C_1$ - $C_6$  alkoxy,  $C_1$ - $C_6$  alkylmercapto,  $C_1$ - $C_6$  alkoxycarbonyl,  $C_1$ - $C_6$  alkylsulfinyl or  $C_1$ - $C_6$  alkylsulfonyl groups;

 $R^2$  is selected from the group consisting of hydrogen, a straight-chain or branched, saturated or unsaturated alkyl chain having 1-20 carbon atoms, which is unsubstituted or substituted at least once by halogen,  $C_1$ - $C_6$  alkylmercapto,  $C_1$ - $C_6$  alkoxycarbonyl or  $C_1$ - $C_6$  alkylsulfonyl groups;

X is selected from the group consisting of a valence bond, an oxygen atom, a sulfur atom, a sulfinyl group and a sulfonyl group;

Y is selected from the group consisting of a valence bond, an oxygen atom and a sulfur atom;

n is an integer of 1, 2 or 3;

whereby the amino group of the nucleoside base may be unsubstituted or substituted by a known amino protecting group, their tautomers, their optically active forms and racemic mixtures, and their physiologically acceptable salts of inorganic and organic acids or bases.

- 21. The nucleotide derivative according to claim 20, wherein  $R^1$  is a straight-chain  $C_8$ - $C_{15}$  alkyl group, which is unsubstituted or substituted by a  $C_1$ - $C_6$  alkoxy or a  $C_1$ - $C_6$  alkylmercapto group.
- 22. The nucleotide derivative according to claim 20, wherein  $R^2$  represents a straight-chain  $C_8$ - $C_{15}$  alkyl group, which is unsubstituted or substituted by a  $C_1$ - $C_6$  alkoxy or a  $C_1$ - $C_6$  alkylmercapto group.
- 23. The nucleotide derivative according to claim 20, wherein X is sulfur, sulfinyl or sulfonyl, and Y is oxygen.
- 24. The nucleotide derivative according to claim 1, wherein X and Y are valence bonds,  $R^2$  is hydrogen, and  $R^1$  is a  $C_1$ - $C_{20}$  alkyl group, which is unsubstituted or substituted by a  $C_1$ - $C_6$  alkoxy or a  $C_1$ - $C_6$  alkylmercapto group.
- 25. The nucleotide derivative according to claims 1 to 6, wherein n is 1.
- 26. The nucleotide derivative according the claim 1 or 20, wherein the compound is:

wherein X is sulfur, sulfinyl or sulfonyl.

- 27. A composition comprising at least one compound according to claim 1 20 in combination with a pharmaceutically acceptable adjuvant or vehicle.
- 28. A method for treating malignant tumors comprising administering to a patient in need of such treatment an amount of the composition according to claim 28 effective to treat said tumors.
- 29. The method according to claim 29, wherein said tumor is selected from the group consisting of carcinomas, sarcomas or leukemias.
- 30. A method for treating malignant tumors comprising administering to a patient in need of such treatment an amount of the composition according to claim 29 effective to treat said tumors in fixed or free combination with other anticancer agents.

## **ABSTRACT**

The subject of the present invention are specific lipidesters of halogenatedadenine nucleotides and the use of such lipidesters in the treatment of tumors.